

The Land Breeze and ClNO_2 production over Santa Monica Bay

Nick Wagner, Steven S. Brown, William P. Dubé, Brian M. Lerner, Eric J. Williams, Wayne Angevine, James Roberts - NOAA Earth System Research Lab, USA
Theran Riedel, and Joel A. Thornton - University of Washington, USA
Derek Coffman - Pacific Marine Environmental Laboratory, NOAA, Seattle, Washington, USA
A. Vlasenko, Shao-Meng Li - Air Quality Research Division, Science and Technology Branch, Environment Canada, Canada

Outline:

- 1) Sea breeze and land breeze in Santa Monica Bay.
- 2) ClNO_2 production in the land breeze.


At
t
l
C
a
L
N
e
x
n
t
i
s



Credit: Dan Fornari @ WHOI

ClNO₂ Production Mechanism


- O₃ and NO₂ react to produce NO₃. NO₃ is photoliable and exists in significant concentrations only at night.
- NO₃ and N₂O₅ are usually in chemical equilibrium and rapidly interconvert.
- NO₃ is lost to reactions with VOCs.
- N₂O₅ is lost to hydrolysis.
- Conventional view: In the aerosol N₂O₅ is converted into NO₃⁻ and HNO₃, which is lost to deposition.
- New mechanism: N₂O₅ is converted to ClNO₂ in aerosol with Cl⁻. ClNO₂ builds throughout the night and is lost to photolysis in the morning.




Cass and Shair Experiment – Observation of the Land Breeze

- 1970's transport study: SF_6 injected into power plant at El Segundo in early evening.
- Inferred transport in land breeze initiated aloft, eventually mixing to surface level.
- Return transport during sea breeze the following day.
- Pre-CalNex view of ClNO_2 production:


ClNO_2 is produced in urban outflow over the ocean. The most consistent outflow is the land breeze over Santa Monica Bay.



F. Shair, Atmos. Environ., 1982



Santa Monica Bay




— R/V Atlantis track ■ Radiosonde Launch
■ Wind Profiler Location

- Use measurements from R/V Atlantis in Santa Monica Bay, LAX wind profiler and P-3 vertical profiles over LAX.


LAX wind profiler

- Located on the coast of Santa Monica Bay.
- Data from mid-April through mid-July.
- Diurnally averaged wind direction shows a land breeze and sea breeze below 700 meters.
- The land breeze occurs on the majority of nights during spring.
- Observations from the R/V Atlantis and the NOAA P-3 were at times consistent with the entire period.








ClNO₂ Observed from the R/V Atlantis

- ClNO₂ is observed in the land breeze at mixing ratios as high as 2 ppbv.
- Use a box model to calculate ClNO₂ production over water and compare with measurements.





Box modeling of ClNO₂ production


• Reactions:



- Wind speed and direction give the transport time from the coast.
- DMS concentration is used to estimate NO₃ loss rate.
- Upper limit for ClNO₂ production, because the maximum N₂O₅ uptake was assumed. ($\gamma = 0.04$)
- Used two different yields:
 - $\phi = 100\%$ for upper limit
 - $\phi = \%$ of NO_z



Example Boxmodel Calculation




- Initialized the model to measured values of O_3 , NO_2 , and N_2O_5 .


Results of Box Modeling

- Four nights in Santa Monica Bay during the land breeze were used.

- ClNO_2 production over water is insufficient to account for observed ClNO_2 .



Aircraft Vertical Profiles over LAX – Transport of ClNO_2 into Santa Monica Bay




- Vertical profiles confirms transport of ClNO_2 into Santa Monica Bay.



Conclusions

Post-CalNex view:

ClNO_2 is produced over Santa Monica Bay. However, significant production occurs over land and some ClNO_2 is transported to Santa Monica Bay on the land breeze.





Production of ClNO_2 over land compared with over water.

- As the aerosol moves inland on the sea breeze it becomes more acidic. The acidification process allows Cl^- to move to particles with the most surface area. Gard, Science 279, 1184 (1998)
- High aerosol liquid water content over water dilutes Cl^- and reduces the yield. (future work?)
- Stronger sea breeze transports more Cl^- in land than NO_x is transported to sea in land breeze.
- See Levi Mielke's poster on ClNO_2 measurements at Pasadena ground site.

Composition of the Land Breeze

- In the land breeze are fresh urban emissions.
- The sea breeze is a more aged and dilute urban air mass.




Details of Boxmodeling


	5/16/2011	5/21/2011	5/24/2011	5/25/2011
Distance to the Coast	6.7 km	12.6 km	2.1 km	7.3 km
Transport time	1.8 hours	2.0 hours	0.18 hours	1.1 hours
N ₂ O ₅	23 ± 6 pptv	49 ± 13 pptv	54 ± 19 pptv	8 ± 3 pptv
NO ₂	9.4 ± 2.1 ppbv	9.0 ± 0.6 ppbv	14.3 ± 3.2 ppbv	3.2 ± 0.8 ppbv
O ₃	35.5 ± 2.4 ppbv	26.4 ± 3.3 ppbv	21.5 ± 2.6 ppbv	34.8 ± 1.0 ppbv
DMS	35 ± 10 pptv	18 ± 11 pptv	21 ± 10 pptv	15 ± 9 ppbv
k _{NO3+DMS}	0.005 s ⁻¹	0.0025 s ⁻¹	0.0035 s ⁻¹	0.0021 s ⁻¹
Total Surface Area	960 ± 50 μm ² /cm ³	1021 ± 50 μm ² /cm ³	550 ± 50 μm ² /cm ³	266 ± 20 μm ² /cm ³
k _{N2O5}	0.0022 s ⁻¹	0.0024 s ⁻¹	0.0013 s ⁻¹	0.00063 s ⁻¹
CINO ₂ produced over water γ = 100%	790 pptv	720 pptv	65 pptv	92 pptv
CINO ₂ produced over water	79 pptv = 10% ϕ	360 pptv = 50% ϕ	32 pptv ϕ = 50%	46 pptv ϕ = 50%
CINO ₂ Observed	320 pptv	1450 ± 250 pptv	580 ± 61 pptv	340 ± 41 pptv

AMS Aerosol Composition: Land Breeze – Sea Breeze


Submicron Aerosol Dry Mass




Aerosol Composition – Filter Samples – Land Breeze




Submicron Land Breeze, filter number 1
5/15 8:19pm - 5/16 7:00 am
mass = 5.6 ug/m3



Submicron Land Breeze, filter number 19
5/23 7:15 pm - 5/24 6:59 am
Mass = 1.3 ug/m3




Supermicron Land Breeze, filter number 1
5/15 8:19pm - 5/16 7:00 am
Mass = 9.4 ug/m3




Supermicron Land Breeze, filter number 19
5/23 7:15 pm - 5/24 6:59 am
Mass = 21.7 ug/m3


Aerosol Composition – Filter Samples – Sea Breeze




Submicron Sea Breeze, filter number 0
5/15 8:11 am - 5/15 1:34 pm
Mass = 10 ug/m3



Submicron Sea Breeze, filter number 15
5/21 8:40 am - 5/21 12:50 pm
Mass = 4.4 ug/m3




Supermicron Sea Breeze, filter number 0
5/15 8:11 am - 5/15 1:34 pm
Mass = 12 ug/m3



Supermicron Sea Breeze, filter number 15
5/21 8:40 am - 5/21 12:50 pm
Mass = 20.9 ug/m3

Radiosondes Launched from the R/V Atlantis


- Four radiosondes launched during the land breeze. (during the night in SMB)
- The marine boundary layer extends up to between 550 and 750 meters.
- Land breeze in the boundary layer from the East - South East.
- Above the marine boundary layer winds are from the northwest.
- R/V Atlantis measurements are representative of the marine boundary layer.



Land Breeze observed on the R/V Atlantis

- The R/V Atlantis observed the land breeze in Santa Monica Bay on five nights. (5/15th, 5/16th, 5/21st, 5/24th and 5/25th)
- The land breeze begins around midnight and continues a few hours after sunrise.

Santa Monica Bay



- R/V Atlantis track
- Radiosonde Launch
- Wind Profiler Location

Diurnal Wind Speed and Direction

